

# On $n$ -sums in an abelian group

Weidong Gao

Center for Combinatorics, LPMC-TJKLC,  
Nankai University, Tianjin 300071, P.R. China

A joint work with

David J. Grynkiewicz and Xingwu Xia

*MSC2010*: Primary 11B75; Secondary 11P99, 20K01

## 1. Introduction

Let  $G$  be an additive abelian group, let  $S$  be a sequence of elements from  $G$ , and let  $|S|$  denote the length of  $S$ . For an integer  $n \geq 1$ , let  $\Sigma_n(S)$  denote the set that consists of all elements in  $G$  which can be expressed as the sum of terms from a subsequence of  $S$  having length  $n$ . The famous Erdős-Ginzburg-Ziv Theorem asserts that, if  $G$  is finite and  $|S| \geq 2|G| - 1$ , then  $0 \in \Sigma_{|G|}(S)$ . This theorem has attracted a lot of attention, and  $\Sigma_{|G|}(S)$  has been studied by many authors.

In 1967, Mann [19] extended this theorem by showing that, if  $|G|$  is prime and every term of  $S$  has multiplicity at most  $|S| - |G| + 1$ , then  $\Sigma_{|G|}(S) = G$ .

In 1977, Olson [21] generalized Mann's result to any finite abelian group and showed that, if  $|S| \geq 2|G| - 1$  and each coset  $x + H$  contains at most  $|S| + 1 - \frac{|G|}{|H|}$  terms of  $S$ , for every subgroup  $H$ , then  $\sum_{|G|}(S) = G$ .

In 1995, the first author [9] proved that Olson's result is true with the restriction  $|S| \geq 2|G| - 1$  replaced by  $|S| \geq |G| + D(G) - 1$ , where  $D(G)$  is the Davenport constant of  $G$ , which is the smallest integer  $d$  such that every sequence over  $G$  of length at least  $d$  has a nonempty zero-sum subsequence.

In 2009, the restriction  $|S| \geq |G| + D(G) - 1$  was further weakened to  $|S| \geq |G| + d^*(G)$  by the second author [17], where  $d^*(G) = \sum_{i=1}^r (n_i - 1)$  when  $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$  with  $n_1 \mid \dots \mid n_r$  (see also [15, Exercise 15.4]). (Note, it is well-known and rather trivial that  $D(G) \geq d^*(G) + 1$ .)

In 1999, Bollobás and Leader [1] proved that, if  $|S| \geq |G| + 1$ , then either  $0 \in \Sigma_{|G|}(S)$  or  $|\Sigma_{|G|}(S)| \geq |S| - |G| + 1$ .

They further conjectured that the minimum of  $|\Sigma_{|G|}(S)|$ , assuming  $0 \notin \Sigma_{|G|}(S)$ , equals the minimum of  $|\Sigma(T)|$ , assuming  $T$  is zero-sum free and  $|T| = |S| - |G| + 1$ , which was confirmed by the first author and Leader [12] in 2005.

In 2003, Y. O. Hamidoune [18] noted that the bounds for  $|\Sigma_{|G|}(S)|$ , assuming  $0 \notin \Sigma_{|G|}(S)$ , seemed to only be tight for sequences having few distinct terms. To make this specific, he made the following two conjectures (for cyclic groups).

**Conjecture 1.1.** *Let  $G$  be a finite abelian group and let  $S$  be a sequence over  $G$  of length  $|S| \geq |G| + 1$ . Suppose the maximum multiplicity of a term of  $S$  is at most  $|G| - |\text{supp}(S)| + 2$ . Then either*

$$|\Sigma_{|G|}(S)| \geq |S| - |G| + |\text{supp}(S)| - 1$$

*or there exists a nontrivial subgroup  $H \leq G$  with  $H \subset \Sigma_{|G|}(S)$ , where  $|\text{supp}(S)|$  denotes the number of distinct terms in  $S$ .*

**Conjecture 1.2.** *Let  $G$  be a finite abelian group and let  $S$  be a sequence over  $G$  of length  $|S| \geq |G| + 1$ . If  $0 \notin \Sigma_{|G|}(S)$ , then*

$$|\Sigma_{|G|}(S)| \geq |S| - |G| + |\text{supp}(S)| - 1,$$

*where  $|\text{supp}(S)|$  denotes the number of distinct terms in  $S$ .*

In 2005, Conjecture 1.1 was resolved by the second author [15]. Later, it was pointed out by DeVos, Goddyn and Mohar [6] that a similar method actually yields the following stronger generalization of Conjecture 1.1.

**Theorem 1.3.** *Let  $G$  be an abelian group, let  $n \geq 1$  be an integer, and let  $S$  be a sequence over  $G$  of length  $|S| \geq n + 1$ . Suppose the maximum multiplicity of a term of  $S$  is at most  $n - |\text{supp}(S)| + 2$ . Then either*

$$|\Sigma_n(S)| \geq \min\{n + 1, |S| - n + |\text{supp}(S)| - 1\}$$

*or there exists a nontrivial subgroup  $H \leq G$  with  $ng + H \subset \Sigma_n(S)$  for some  $g \in \text{supp}(S)$ , where  $|\text{supp}(S)|$  denotes the number of distinct terms in  $S$ .*

In this paper, we show the following similar result to Theorem 1.3 and confirm Conjecture 1.2 as its corollary.

**Theorem 1.4.** *Let  $G$  be an abelian group, let  $n \geq 1$  be an integer, let  $S$  be a sequence over  $G$  of length  $|S| \geq n + 1$ , and let  $h(S)$  denote the maximum multiplicity of a term from  $S$ . Then either*

*$|\Sigma_n(S)| \geq \min\{n + 1, |S| - n + |\text{supp}(S)| - 1\}$   
or  $ng \in \Sigma_n(S)$  for every  $g \in G$  whose multiplicity in  $S$  is at least  $v_g(S) \geq h(S) - 1$ , where  $|\text{supp}(S)|$  denotes the number of distinct terms in  $S$ .*

Taking  $G$  finite and  $n = |G|$  in the above theorem, Conjecture 1.2 clearly follows. For some related papers, we refer to [2, 3, 5, 8, 10, 11, 20, 21, 24].

## 2. Notation and Preliminaries

Let  $\mathbb{N}$  denote the set of positive integers and let  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . For any two integers  $a, b \in \mathbb{N}_0$ , we set  $[a, b] = \{x \in \mathbb{N}_0 : a \leq x \leq b\}$ . Throughout this paper, all abelian groups will be written additively.

Let  $G$  be an abelian group and let  $\mathcal{F}(G)$  be the free abelian monoid, multiplicatively written, with basis  $G$ . The elements of  $\mathcal{F}(G)$  are simply finite (unordered) sequences with terms from  $G$ , multiplicatively written. We write sequences  $S \in \mathcal{F}(G)$  in the form

$$S = \prod_{g \in G} g^{v_g(S)}, \text{ with } v_g(S) \in \mathbb{N}_0 \text{ for all } g \in G.$$

We call  $v_g(S)$  the multiplicity of the term  $g$  in  $S$  and say that  $S$  contains  $g$  if  $v_g(S) > 0$ . Furthermore,  $S$  is called square-free if  $v_g(S) \leq 1$  for all  $g \in G$ . The unit element  $1 \in \mathcal{F}(G)$  is called the empty sequence. We use  $S_1 \mid S$  to denote that the sequence  $S_1$  is a subsequence

of  $S$ . In such case,  $SS_1^{-1}$  denotes the subsequence of  $S$  obtained by removing the terms from  $S_1$ . Let  $S_1, \dots, S_r$  be subsequences of  $S$ . We say  $S_1, \dots, S_r$  are disjoint subsequences if  $S_1 \cdot \dots \cdot S_r \mid S$ . If a sequence  $S \in \mathcal{F}(G)$  is written in the form  $S = g_1 \cdot \dots \cdot g_\ell$ , we tacitly assume that  $\ell \in \mathbb{N}_0$  and  $g_1, \dots, g_\ell \in G$ .

For a sequence

$$S = g_1 \cdot \dots \cdot g_\ell = \prod_{g \in G} g^{v_g(S)} \in \mathcal{F}(G),$$

we call

- $|S| = \ell = \sum_{g \in G} v_g(S) \in \mathbb{N}_0$  the *length* of  $S$ ,
- $h(S) = \max\{v_g(S) : g \in G\} \in [0, |S|]$  the *maximum* of the *multiplicities* of  $S$ ,
- $\text{supp}(S) = \{g \in G : v_g(S) > 0\} \subset G$  the *support* of  $S$ ,
- $\sigma(S) = \sum_{i=1}^{\ell} g_i = \sum_{g \in G} v_g(S)g \in G$  the *sum* of  $S$ .

If  $\phi : G \rightarrow G'$  is a map, then  $\phi(S) = \phi(g_1) \cdot \dots \cdot \phi(g_\ell) \in \mathcal{F}(G')$  denotes the sequence over

$G'$  obtained by applying  $\phi$  to each term of  $S$ .  
Note  $|\phi(S)| = |S|$ .

For  $r \in \mathbb{Z}$ , we define

$$\Sigma_r(S) = \{\sigma(S') : S' \mid S \text{ and } |S'| = r\}.$$

Note  $\sigma(S') = 0$  when  $S'$  is the empty sequence.

For  $k \in \mathbb{Z}$ , define

$$\Sigma_{\geq k}(S) = \bigcup_{r=k}^{\ell} \Sigma_r(S), \quad \Sigma_{\leq k}(S) = \bigcup_{r=1}^k \Sigma_r(S) \quad \text{and} \quad \Sigma$$

and

$$\Sigma_{\leq k}^*(S) = \{0\} \cup \Sigma_{\leq k}(S) \quad \text{and} \quad \Sigma^*(S) = \{0\} \cup \Sigma(S).$$

A sequence  $S$  is called

- a *zero-sum sequence* if  $\sigma(S) = 0$ ,
- *zero-sum free* if  $0 \notin \Sigma(S)$ .

Let  $A$  and  $B$  be two nonempty subsets of  $G$ .  
Define

$$A + B = \{a + b : a \in A, b \in B\}.$$

If  $A = \{x\}$  for some  $x \in G$ , then we simply denote  $A + B$  by  $x + B$ . For any nonempty subset  $C$  of  $G$ , let  $-C = \{-c : c \in C\}$ . We say that

$g \in G$  is a unique expression element of  $A + B$  if there is precisely one pair  $(a, b) \in A \times B$  with  $a + b = g$ . For a nonempty subset  $A \subset G$  and a subgroup  $H$  of  $G$ , we say that  $A$  is  $H$ -periodic if  $A$  is a union of  $H$ -cosets. Let  $\text{stab}(A)$  denote the stabilizer of  $A$  in  $G$ , i.e.,  $\text{stab}(A) = \{g \in G : g + A = A\}$ . Then  $\text{stab}(A)$  is the maximal subgroup  $H$  for which  $A$  is  $H$ -periodic. The set  $A$  is called *periodic* if  $\text{stab}(A)$  is nontrivial. We use  $\phi_H : G \rightarrow G/H$  for the natural homomorphism.

To prove Theorem 1.4, we need some preliminaries, beginning with a result of Scherk [25].

**Lemma 2.1.** *Let  $G$  be an abelian group and let  $A$  and  $B$  be two finite subsets of  $G$  such that  $A + B$  contains a unique expression element. Then  $|A + B| \geq |A| + |B| - 1$ .*

By using Lemma 2.1 repeatedly, one can prove the following result of Bovey, Erdős and Niven [4].

**Lemma 2.2.** *Let  $S$  be a zero-sum free sequence over an abelian group and let  $S_1, \dots, S_k$  be disjoint subsequences of  $S$ . Then*

$$|\Sigma(S)| \geq \sum_{i=1}^k |\Sigma(S_i)| \quad \text{with} \quad |\Sigma(S_i)| \geq |S_i| \quad \text{for all } i.$$

We also need the following result, which is the common corollary of two more general additive results: the DeVos-Goddyn-Mohar Theorem and the Partition Theorem (see [16, Chapters 13-14]).

**Theorem 2.3.** [6, 16] *Let  $G$  be an abelian group. If  $S$  is a sequence over  $G$ ,  $n \leq |S|$ , and  $H = \text{stab}(\Sigma_n(S))$ , then*

$$|\Sigma_n(S)| \geq \left( \sum_{g \in G/H} \min\{n, v_g(\phi_H(S))\} - n + 1 \right) |H|,$$

where  $v_g(\phi_H(S))$  denotes the multiplicity of the term  $g \in G/H$  in the sequence  $S$  when its terms have been reduced modulo  $H$ .

**Lemma 2.4.** *Let  $G$  be an abelian group, let  $n \geq 1$  be an integer, let  $S \in \mathcal{F}(G)$  be a sequence over  $G$  with*

$$|\Sigma_n(S)| \leq |S| - n,$$

*let  $H = \text{stab}(\Sigma_n(S))$ , and let  $\phi_H : G \rightarrow G/H$  be the natural homomorphism.*

*1. If  $h(S) \leq n$  and  $g \in \text{supp}(S)$  is a term with  $v_{\phi_H(g)}(\phi_H(S)) \geq n$ , then*

$$v_{\phi_H(g)}(\phi_H(S)) \geq n + |H|.$$

*2. If  $g \in G$  is a term with near maximum multiplicity  $v_g(S) \geq h(S) - 1$ , then*

$$v_{\phi_H(g)}(\phi_H(S)) \geq n.$$

*Moreover, the above inequality is strict if either  $h(S) \leq n$  or  $v_g(S) = h(S)$ .*

*Proof.* Observe that  $0 \leq |\Sigma_n(S)| \leq |S| - n$  implies  $|S| \geq n$ . Applying Theorem 2.3 to  $\Sigma_n(S)$ , we

find that

$$(1) \quad |\Sigma_n(S)| \geq \left( \sum_{g \in G/H} \min\{n, v_g(\phi_H(S))\} - n + 1 \right) |H|.$$

Let  $N \geq 0$  denote the number of  $g \in G/H$  with  $v_g(\phi_H(S)) \geq n$  and let  $e$  denote the number of terms of  $S$  not equal modulo  $H$  to some  $g \in G/H$  with  $v_g(\phi_H(S)) \geq n$ . Then (1) can be rewritten as

$$(2) \quad |\Sigma_n(S)| \geq ((N - 1)n + e + 1)|H|,$$

and we clearly have

$$(3) \quad |S| \leq h(S)N|H| + e.$$

If  $N = 0$ , then  $e = |S|$ , whence (2) yields  $|\Sigma_n(S)| \geq (|S| - n + 1)|H| \geq |S| - n + 1$ , contrary to hypothesis. Therefore we may assume

$$N \geq 1.$$

Combining (2), (3) and the hypothesis  $|\Sigma_n(S)| \leq |S| - n$  yields

(4)

$$((N-1)n+e+1)|H| \leq |\Sigma_n(S)| \leq |S|-n \leq h(S)N|H|+e-n.$$

1. Let  $x = v_{\phi_H(g)}(\phi_H(S))$ . Then, since  $v_{\phi_H(g)}(\phi_H(S)) \geq n$ , we can improve (3) to

$$|S| \leq h(S)(N-1)|H| + e + x.$$

Thus we can improve (4) to

$$((N-1)n+e+1)|H| \leq |\Sigma_n(S)| \leq |S|-n \leq h(S)(N-1)|H|+e+x$$

which rearranges to give

$$v_{\phi_H(g)}(\phi_H(S)) = x \geq (N-1)|H|(n-h(S))+e(|H|-1)+n+1$$

Since  $h(S) \leq n$ , applying the estimates  $N \geq 1$  and  $e \geq 0$  yields the desired lower bound.

2. If the second conclusion of this lemma is false, then every term of  $S$  equal to  $g$  is counted by  $e$ , i.e.,

$$e \geq v_g(S) \geq h(S) - 1.$$

Rearranging (4) and applying the above estimate, we obtain

$$\begin{aligned}
0 &\geq (n - h(S))N|H| + e(|H| - 1) - n(|H| - 1) + |H| \\
&\geq (n - h(S))N|H| + (h(S) - 1)(|H| - 1) - n(|H| - 1) \\
&= (n - h(S))(N|H| - |H| + 1) + 1.
\end{aligned}$$

Hence, since  $N \geq 1$ , it follows that  $h(S) \geq n + 1$ , in which case  $v_{\phi_H(g)}(\phi_H(S)) \geq v_g(S) \geq h(S) - 1 \geq n$ , a contradiction.

If  $h(S) \leq n$ , then part 1 now implies  $v_{\phi_H(g)}(\phi_H(S)) \geq n + |H| \geq n + 1$ . On the other hand, if  $h(S) \geq n + 1$  and  $v_g(S) = h(S)$ , then we trivially have  $v_{\phi_H(g)}(\phi_H(S)) \geq v_g(S) = h(S) \geq n + 1$ , completing the proof.  $\square$

The following lemma is crucial in this paper.

**Lemma 2.5.** *Let  $G$  be an abelian group, let  $n \geq \lambda \geq 0$  be integers, and let  $S = T0^{n-\lambda} \in \mathcal{F}(G)$  be a sequence over  $G$  with  $|S| \geq n$  and  $v_0(S) \geq h(S) - 1$ . Then either  $|\Sigma_n(S)| \geq n + 1$  or*

$$\Sigma_{\geq \lambda}(T) = \Sigma_n(S).$$

*Proof.* Observe that

$$\Sigma_n(S) = \Sigma_n(T0^{n-\lambda}) = \Sigma_{[\lambda, n]}(T) = \{\sigma(T') : T' \mid T \text{ and}$$

Thus  $\Sigma_{\geq \lambda}(T) = \Sigma_n(S)$  is trivial unless

$$|T| \geq n + 1,$$

which we now assume. This also shows that  $\Sigma_n(S) \subset \Sigma_{\geq \lambda}(T)$ , so that it suffices to show  $\Sigma_{\geq \lambda}(T) \subset \Sigma_n(S)$ . Moreover, we have  $|S| \geq |T| \geq n + 1 \geq \lambda + 1$ , so that  $|T| - \lambda \geq 1$ .

Now

$$\Sigma_n(S) = \sigma(S) - \Sigma_{|S|-n}(S) = \sigma(T) - \Sigma_{|T|-\lambda}(S) \quad \text{and}$$

Thus to show  $\Sigma_{\geq \lambda}(T) \subset \Sigma_n(S)$ , it suffices to show

$$(5) \quad \Sigma_{\leq |T|-\lambda}^*(T) \subset \Sigma_{|T|-\lambda}(S),$$

and to show  $|\Sigma_n(S)| \geq n + 1$ , it suffices to show  $|\Sigma_{|T|-\lambda}(S)| \geq n + 1$ . We now assume

$$(6) \quad |\Sigma_{|T|-\lambda}(S)| \leq n = |S| - (|T| - \lambda)$$

and proceed to establish (5).

Let  $H \leq G$  denote the stabilizer of  $\Sigma_{|T|-\lambda}(S)$ . Then, in view of (6) and the hypothesis  $v_0(S) \geq h(S) - 1$ , we can apply Lemma 2.4.2 to conclude that

$$(7) \quad v_0(\phi_H(S)) \geq |T| - \lambda.$$

In particular,  $\phi_H(T_{G \setminus H})0^{|T|-\lambda}$  is a subsequence of  $\phi_H(S)$ , where  $T_{G \setminus H} | T$  denotes the subsequence consisting of all terms from  $G \setminus H$ . Consequently, since  $\Sigma_{|T|-\lambda}(S)$  is  $H$ -periodic, we see that, in order to establish (5) (and thus complete the proof), it suffices to show

$$\Sigma_{\leq |T|-\lambda}^*(\phi_H(T_{G \setminus H})) = \Sigma_{\leq |T|-\lambda}^*(\phi_H(T)) \subset \Sigma_{|T|-\lambda}(\phi_H(T_C))$$

Since the above inclusion holds trivially with equality, the proof is complete.  $\square$

If  $A \subset G$ , then we define  $\Sigma(A) = \Sigma(S)$  where  $S$  is the square-free sequence with  $\text{supp}(S) = A$ .

**Lemma 2.6.** *Let  $S$  be a subset of an abelian group  $G$  with  $0 \notin \Sigma(S)$ . Then*

- (1)  $|\Sigma(S)| \geq 2|S| - 1$ ;
- (2) if  $|S| \geq 4$ , then  $|\Sigma(S)| \geq 2|S|$ ;
- (3) if  $|S| = 3$  and  $S$  does not contain exactly one element of order two, then  $|\Sigma(S)| \geq 2|S|$ .

*Proof.* 1. and 2. have been proved in [7].

3. If  $S$  contains no element of order two, then the result has also been proved in [7]. Now assume that  $S$  contains at least two elements of order two. Let  $S = \{a, b, c\}$  with  $\text{ord}(a) = \text{ord}(b) = 2$ . If  $c = a + b$ , then  $a + b + c = a + b + a + b = 2a + 2b = 0 + 0 = 0$ , contradicting that  $0 \notin \Sigma(S)$ . Therefore,  $a + b \notin S$ . If  $a + c = b$ , then  $a + c + b = 2b = 0$ , likewise a contradiction. Hence,  $a + c \notin S$ . Similarly, we can prove  $b + c \notin S$ . Note that  $a + b + c \notin$

$\{a, b, c, a + b, b + c, c + a\}$ . Therefore,  $|\Sigma(S)| = 7$  and we are done.  $\square$

**Lemma 2.7.** *Let  $G$  be an abelian group and let  $S \in \mathcal{F}(G)$  be a zero-sumfree sequence. Then  $|\Sigma(S)| \geq |S| + |\text{supp}(S)| - 1$ , and we have strict inequality unless  $|S| \leq 2$  or  $|S| = 3$  with  $S$  containing exactly one element of order two.*

*Proof.* Let  $S_1$  be a square-free subsequence of  $S$  with  $|S_1| = |\text{supp}(S)|$  and let  $S_2 = S S_1^{-1}$ . Applying Lemma 2.2 to  $S = S_1 S_2$ , we obtain that

$$|\Sigma(S)| \geq |\Sigma(S_1)| + |\Sigma(S_2)| \geq |S_2| + |\Sigma(S_1)| = |S| - |S_1| + |\Sigma(S_1)|$$

Now the result follows from Lemma 2.6.  $\square$

Given subsets  $A, B \subset G$ , we define the restricted sumset to be

$$A \dot{+} B = \{a + b : a \in A, b \in B, a \neq b\}.$$

**Lemma 2.8.** *Let  $A$  be a finite subset of an abelian group with  $0 \in A$  and  $|A| \geq 3$  and let  $H = \langle A \rangle$ . If*

*H is an elementary 2-group, also suppose that  $A \neq H$ . Then  $|A+A| \geq |A|$ .*

*Proof.* Assume by contradiction that  $|A+A| \leq |A| - 1$ . Clearly,  $a + A \setminus \{a\} \subset A+A$  for all  $a \in A$ . Thus

$$(8) \quad a + A \setminus \{a\} = A+A = A \setminus \{0\}$$

for all  $a \in A$ .

If every nonzero element of  $A$  has order 2, then  $H$  will be an elementary 2-group and  $A+A = (A + A) \setminus \{0\}$ . In this case, (8) implies  $A = A + A$ , which is easily seen to only be possible if  $A$  is itself a subgroup, thus equal to  $H$ . As this is contrary to hypothesis, we may now assume there is some  $a \in A \setminus \{0\}$  with  $\text{ord}(a) \geq 3$ .

Now (8) is only possible if

$$A = \{0, a\} \cup B$$

with  $B = a + B$  a disjoint  $\langle a \rangle$ -periodic subset. Since  $\langle a \rangle$  is a cyclic group of order at least 3, and since  $B$  is  $\langle a \rangle$ -periodic, it follows that

$B \dot{+} B = B + B \subset A \dot{+} A = \{a\} \cup B$  is also  $\langle a \rangle$ -periodic. Thus  $B + B = B$ , which is only possible if  $B$  is a subgroup of  $G$  or the empty set. Since  $0 \notin B$ , the former is not possible, and since  $|A| \geq 3$ , the latter is also not possible, a concluding contradiction.  $\square$

**Lemma 2.9.** *Let  $A$  be a finite subset of an abelian group with  $0 \in A$  and  $|A| \geq 4$  and let  $H = \langle A \rangle$ . Suppose  $|A| \leq |H| - 1$  with strict inequality if  $H$  is an elementary 2-group. Then  $|A \dot{+} A| \geq |A| + 1$  or  $A = L \cup (a + L)$  for some cardinality two subgroup  $L \leq G$  and  $a \in G$ .*

*Proof.* Assume by contradiction that  $|A \dot{+} A| \leq |A|$ . By Lemma 2.8, we have

$$|A \dot{+} A| = |A|.$$

Clearly,  $a + A \setminus \{a\} \subset A \dot{+} A$  for all  $a \in A$ . Thus

$$(9) \quad a + A \setminus \{a\} \subset A \dot{+} A = (A \setminus \{0\}) \cup \{b\}$$

for all  $a \in A$  and some  $b \notin A \setminus \{0\}$ .

If every nonzero element of  $A$  has order 2, then  $H$  will be an elementary 2-group and  $A \dot{+} A =$

$(A + A) \setminus \{0\}$ . In this case, (9) implies  $A + A = A \cup \{b\}$ , which, in view of  $|A| \geq 3$ , is only possible if  $A$  is itself a subgroup or a subgroup with at most one element removed (being a simple consequence of Kneser's Theorem [16, Chapter 6]). Hence  $|A| \geq |H| - 1$ , contrary to hypothesis, and we may now assume there is some  $a \in A \setminus \{0\}$  with  $\text{ord}(a) \geq 3$ . Let  $K = \langle a \rangle$ .

Now (9) is only possible if

$$A = \{0, a\} \cup B \cup B'$$

with  $B = B + a$  a disjoint  $K$ -periodic subset and  $B'$  either empty or a disjoint arithmetic progression with difference  $a$  whose last term is  $b - a$ . Since  $\text{ord}(a) \geq 3$ ,  $K$  is a cyclic group of order at least 3.

Suppose  $B$  is nonempty. Then, since  $B$  is  $K$ -periodic with  $K$  a cyclic group of order  $|K| \geq 3$ , it follows that  $A + B = A \dot{+} B \subset A \dot{+} A = (A \setminus \{0\}) \cup \{b\}$ . Since  $A + B$  is  $K$ -periodic, it must be contained in the maximal  $K$ -periodic subset of

$(A \setminus \{0\}) \cup \{b\}$ . We consider two cases depending on whether  $b = 0$  or  $b \neq 0$ .

If  $b = 0$ , then  $(A \setminus \{0\}) \cup \{b\} = A$ . In this case, since  $|\phi_K(A+B)| \geq |\phi_K(A)|$ , we see that the only way  $A + B$  can be contained in the maximal  $K$ -periodic subset of  $A = (A \setminus \{0\}) \cup \{b\}$  is if  $A$  is itself  $K$ -periodic with  $K$  cyclic of order  $|K| \geq 3$ . It follows that  $A+A = A+A = (A \setminus \{0\}) \cup \{b\} = A$ , implying that  $A$  is itself a subgroup, thus equal to  $H$ , which is contrary to hypothesis.

If  $b \neq 0$ , then  $0, a \in A \cap K$  ensures that  $K$  is a  $K$ -coset that intersects  $(A \setminus \{0\}) \cup \{b\}$  but which is not contained in  $(A \setminus \{0\}) \cup \{b\}$ . Consequently, the maximal  $K$ -periodic subset of  $(A \setminus \{0\}) \cup \{b\}$  is contained in  $(A + K) \setminus K$ , and thus has size at most  $|\phi_K(A)| - 1$ . But this makes it impossible for  $A + B$  to be contained in this maximal  $K$ -periodic subset in view of  $|\phi_K(A+B)| \geq |\phi_K(A)|$ . So we may now assume  $B$  is empty.

Since  $B$  is empty and  $|A| \geq 4$ , we have

$$A = \{0, a\} \cup B' = \{0, a\} \cup \{x, x + a, \dots, x + ta\},$$

for some  $x \in G$ , where  $t = |A| - 3 \geq 1$  and  $b = x + (t + 1)a$ . Thus

$$\begin{aligned} A+A &= \{a\} \cup \{x, x + a, \dots, x + (t+1)a\} \cup \{2x + a, 2x + 2a, \dots, 2x + ta\} \\ &= \{a\} \cup \{x, x + a, \dots, x + (t+1)a\}, \end{aligned}$$

with the latter equality from (9) and the elements listed in (11) distinct.

Since  $1 \leq t \leq 2t-1$ , it follows that the element  $2x+ta$ , from the third set in (10), must also lie in the set  $\{a\} \cup \{x, x + a, \dots, x + (t+1)a\}$  from (11). If  $2x + ta = x + ja$  for some  $j \in [0, t]$ , then  $0 = x + (t-j)a \in \{x, x + a, \dots, x + ta\}$ , contradicting that these are all elements of  $A$  distinct from 0 and  $a$ . If  $2x + ta = x + (t+1)a$ , then this implies  $x = a$ , contradicting that  $x, a \in A$  are distinct elements of  $A$ . Therefore the only remaining possibility is that

$$(12) \quad 2x + ta = a.$$

Suppose  $|A| \geq 5$ , which is equivalent to assuming  $t \geq 2$ . In this case, (10) and (12) ensure that  $2a = 2x + (t + 1)a \in A \dot{+} A$ . Comparing this with (11), we see that  $2a \in A \dot{+} A$  forces  $x = 2a$ , which combined with (12) yields  $(t + 3)a = 0$ . Since  $x = 2a$  and  $(t + 3)a = 0$ , it follows that  $A = \{0, a, x, x + a, \dots, x + ta\} = \{0, a, 2a, \dots, (t + 2)a\} = H$ , contrary to hypothesis. So it only remains to consider the case  $|A| = 4$ .

For  $|A| = 4$ , we have  $A = \{0, a\} \cup \{x, x + a\}$ . In this case,

$$A \dot{+} A = \{a\} \cup \{x, x + a, x + 2a\} \cup \{2x + a\}.$$

Since  $A = \{0, a\} \cup \{x, x + a\}$  are the distinct elements of  $A$  with  $\text{ord}(a) \geq 3$ , it is easily verified that the elements  $\{x, x + a, x + 2a\}$  are distinct from each other as well as from  $a$  and  $2x + a$ . Thus  $|A \dot{+} A| \geq 5 = |A| + 1$  follows unless  $a = 2x + a$ . However, if  $a = 2x + a$ , then  $A = \{0, x\} \cup (a + \{0, x\})$  with  $\{0, x\} = L \leq G$  a subgroup of order two, also as desired.  $\square$

Note that Lemmas 2.8 and 2.9 both may be paraphrased as concluding that either  $|A+A|$  is large or  $A$  is a large subset of a periodic subset. Unlike the case of ordinary sumsets, this latter conclusion does not force  $A+A$  to be itself periodic. As yet, there is no Kneser-type extension of the Erdős-Heilbronn Conjecture to an arbitrary abelian group (see [16, Chapter 22]). Lemmas 2.8 and 2.9 may be viewed as the first easily verified cases in whatever this extension should be.

### 3. Proof of Theorem 1.4

*Proof of Theorem 1.4.* Assume by contradiction that we have some  $g \in G$  with  $v_g(S) \geq h(S) - 1$  and  $ng \notin \Sigma_n(S)$ . Note that this theorem is translation invariant, so we may assume that  $g = 0$ . Hence

$$0 = n0 \notin \Sigma_n(S) \quad \text{and} \quad v_0(S) \geq h(S) - 1.$$

If  $v_0(S) \geq n$ , then  $0 = n0 \in \Sigma_n(S)$  holds trivially, contrary to assumption. So we may assume that

$$v_0(S) = n - \lambda \quad \text{for some } \lambda \in [1, n].$$

Let

$$S = 0^{n-\lambda}T$$

with  $0 \nmid T$ . We need to show

$$|\Sigma_n(S)| \geq \min\{n + 1, |S| - n + |\text{supp}(S)| - 1\}.$$

Assume by contradiction that

$$|\Sigma_n(S)| \leq n.$$

Then, by Lemma 2.5,

$$(13) \quad \Sigma_{\geq \lambda}(T) = \Sigma_n(S).$$

So it suffices to prove that

$$|\Sigma_{\geq \lambda}(T)| \geq |S| - n + |\text{supp}(S)| - 1.$$

Let  $T_0$  be a maximal (in length) subsequence of  $T$  with  $\sigma(T_0) = 0$  ( $T_0$  is the empty sequence if  $T$  is zero-sum free). Since  $0 \notin \Sigma_n(S) = \Sigma_{\geq \lambda}(T)$ , we have

$$|T_0| \leq \lambda - 1.$$

Let  $T_1 = TT_0^{-1}$ , so

(14)

$$T = T_0T_1 \quad \text{with} \quad |T_1| = |T| - |T_0| \geq |T| - \lambda + 1 = |S| - n$$

Then, in view of the maximality of  $T_0$ , it follows that

$T_1$  is zero-sum free.

**Claim 1.**  $(\text{supp}(T_0) \setminus \text{supp}(T_1)) \cap \Sigma(T_1) = \emptyset$ .

Assume to the contrary that  $x = \sigma(V_1) \in \text{supp}(T_0) \setminus \text{supp}(T_1)$  for some nontrivial subsequence  $V_1 \mid T_1$ . Then  $|V_1| \geq 2$  (else  $x \in \text{supp}(T_1)$ , contrary to assumption). Therefore,  $T_0x^{-1}V_1$  is a zero-sum subsequence of  $T$  of length  $|T_0| - 1 + |V_1| > |T_0|$ , contradicting the maximality of  $T_0$ . This proves Claim 1.

In view of (14) and the hypothesis  $|S| \geq n + 1$ , choose a subsequence  $V$  of  $T_1$  with

$$(15) \quad |V| = |S| - n - 1$$

and let  $U = T_1 V^{-1}$ . Observe that  $|U| = |T_1| - |V| = |T| - |T_0| - (|S| - n - 1) = \lambda - |T_0| + 1$ , so

$$(16) \quad T_1 = UV \quad \text{with} \quad |U| = \lambda - |T_0| + 1 \geq 2.$$

Furthermore, choose  $V$  as above so that  $|\text{supp}(V) \cap \text{supp}(U)|$  is maximal.

Let

$$A = \{0\} \cup -(\text{supp}(T_0) \setminus \text{supp}(T_1)).$$

Since  $\sigma(T_0) = 0$ , we have

$$(17) \quad A \subset \{0\} \cup -\text{supp}(T_0) = \Sigma_{\geq |T_0|-1}(T_0).$$

Let

$$B = \sigma(U) + \Sigma^*(V).$$

Since  $UV = T_1$ , (16) implies that

$$(18) \quad B \subset \Sigma_{\geq \lambda - |T_0| + 1}(T_1).$$

Since  $T_0 \mid T$  with  $0 \nmid T$ , and since  $V \mid T_1$  with  $T_1$  zero-sum free, we clearly have

$$(19)$$

$$|A| = |\text{supp}(T_0) \setminus \text{supp}(T_1)| + 1 \quad \text{and} \quad |B| = 1 + |\Sigma(V)|$$

Since  $T = T_0 T_1$ , (17) and (18) imply that

$$(20) \quad A + B \subset \Sigma_{\geq \lambda}(T).$$

Let

$$C = \Sigma_{|U|-1}(U) = \sigma(U) - \text{supp}(U).$$

Then

$$(21) \quad |C| = |\text{supp}(U)|.$$

For any  $x \in C$ , there is some subsequence  $U_x \mid U$  with

$$\sigma(U_x) = x \quad \text{and} \quad |U_x| = |U| - 1 = \lambda - |T_0|.$$

Since  $\sigma(T_0) = 0$ , it follows that  $\sigma(U_x T_0) = \sigma(U_x) + \sigma(T_0) = x$  with  $|U_x T_0| = |U_x| + |T_0| = \lambda$ . Since  $U_x \mid U$ ,  $U \mid T_1$  and  $T = T_1 T_0$ , it follows that  $U_x T_0 \mid T$ . As this is true for any  $x \in C$ , we conclude that

$$(22) \quad C \subset \Sigma_\lambda(T) \subset \Sigma_{\geq \lambda}(T).$$

**Claim 2.**  $|A + B| \geq |A| + |B| - 1$ .

Since  $0 \in A$  and  $\sigma(U) \in B$ , we have  $\sigma(U) \in A + B$ . If  $\sigma(U)$  is not a unique expression element of  $A + B$ , then we deduce that  $\sigma(U) = -x + \sigma(U) + \sigma(V_1)$  for some  $x \in \text{supp}(T_0) \setminus \text{supp}(T_1)$  and some nontrivial subsequence  $V_1$  of  $V \mid T_1$ .

It follows that  $\sigma(V_1) = x$ , contrary to Claim 1. Therefore,  $\sigma(U)$  is a unique expression element of  $A + B$ , and Claim 2 follows from Lemma 2.1.

**Claim 3.**  $(A + B) \cap C = \emptyset$ .

Assume to the contrary that Claim 3 is false. We have the following possibilities:

(a)  $\sigma(U) - x = \sigma(U) + \sigma(V_1)$  with  $x \in \text{supp}(U)$  and  $V_1 \mid V$ ; or

(b)  $\sigma(U) - x = \sigma(U) - z + \sigma(V_1)$  with  $x \in \text{supp}(U)$ ,  $z \in \text{supp}(T_0) \setminus \text{supp}(T_1)$  and  $V_1 \mid V$ .

Possibility (a) implies that  $\sigma(xV_1) = 0$ . Since  $V_1 \mid V$ ,  $T_1 = UV$  and  $x \in \text{supp}(U)$ , we must have  $xV_1 \mid T_1$ . But this contradicts that  $T_1$  is zero-sum free. Possibility (b) implies that  $\sigma(xV_1) = z \in \text{supp}(T_0) \setminus \text{supp}(T_1)$ . As before,  $xV_1 \mid T_1$ , and now we have a contradiction to Claim 1. This proves Claim 3.

Now, from (20), (22) and Claim 3, (21), Claim 2, (19), Lemma 2.7 applied to  $\Sigma(V)$  (note  $V \mid T_1$

with  $T_1$  zero-sum free, so  $V$  is also zero-sum free), (15) and the inclusion-exclusion principle,  $T_1 = UV$ ,  $T = T_1T_0$ ,  $\text{supp}(S) \setminus \{0\} \subset \text{supp}(T)$  (which follows from the definition of  $T$ ), and the trivial estimate  $|\text{supp}(U) \cap \text{supp}(V)| \geq 0$ , we obtain

$$\begin{aligned}
|\Sigma_{\geq \lambda}(T)| &\geq |A + B| + |C| \\
&= |A + B| + |\text{supp}(U)| \\
&\geq |A| + |B| - 1 + |\text{supp}(U)| \\
&= |\text{supp}(T_0) \setminus \text{supp}(T_1)| + 1 + |\Sigma(V)| + |\text{supp}(V)| \\
&\geq |\text{supp}(T_0) \setminus \text{supp}(T_1)| + |V| + |\text{supp}(V)| + |\text{supp}(V)| \\
&= |\text{supp}(T_0) \setminus \text{supp}(T_1)| + |S| - n - 1 + |\text{supp}(V)| \\
&= |S| - n - 1 + |\text{supp}(T_0) \setminus \text{supp}(T_1)| + |\text{supp}(V)| \\
&= |S| - n - 1 + |\text{supp}(T)| + |\text{supp}(U) \cap \text{supp}(V)| \\
&\geq |S| - n - 2 + |\text{supp}(S)| + |\text{supp}(U) \cap \text{supp}(V)| \\
&\geq |S| - n - 2 + |\text{supp}(S)|.
\end{aligned}$$

If  $|\Sigma_{\geq \lambda}(T)| \geq |S| - n + |\text{supp}(S)| - 1$ , then the proof is complete. Otherwise, it forces equality in all estimates used above. In particular,

(23)

$$\text{supp}(U) \cap \text{supp}(V) = \emptyset \quad \text{and} \quad |\Sigma(V)| = |V| + |\text{supp}(V)|$$

Now  $\text{supp}(U) \cap \text{supp}(V) = \emptyset$  is only possible, in view of the maximality of  $|\text{supp}(U) \cap \text{supp}(V)|$ , if

$V$  is the empty sequence or  $T_1 = UV$  is square-free.

If  $V$  is empty, then (15) gives  $|S| = n + |V| + 1 = n + 1$ . Clearly,

$$|\Sigma_n(S)| = |\Sigma_{|S|-1}(S)| = |\sigma(S) - \text{supp}(S)| = |\text{supp}(S)| = |S|$$

and we are done. So we may instead assume

$$|V| \geq 1 \quad \text{and} \quad T_1 = UV \quad \text{is square-free.}$$

The estimate  $|\Sigma(V)| = |V| + |\text{supp}(V)| - 1$  from (23) can only hold, according to Lemma 2.7, if

$$(24) \quad |S| - n - 1 = |V| \leq 3,$$

where the first equality follows from (15). This gives us three remaining cases based on the size of  $|V| \in [1, 3]$ .

If  $|V| = |S| - n - 1 = 3$ , then (14) ensures that  $|T_1| \geq |S| - n + 1 = 5$ . Consequently, since  $T_1 = UV$  is square-free, we can choose  $V$  such that  $V$  either contains no element with order two or at least two elements with order two (while still

preserving that  $|\text{supp}(V) \cap \text{supp}(U)| = 0$  is maximal for the definition of  $U$  and  $V$ ). But now Lemma 2.7 ensures that  $|\Sigma(V)| \geq |V| + |\text{supp}(V)|$ , contrary to (23). Therefore it remains to consider the cases when

$$(25) \quad 2 \leq |V| + 1 = |S| - n \leq 3.$$

Note that  $|\Sigma_{\geq \lambda}(T)| = |\sigma(T) - \Sigma_{\leq |T| - \lambda}^*(T)| = |\Sigma_{\leq |T| - \lambda}^*(T)| = |\{0\} \cup \Sigma_{\leq |S| - n}(T)|$  with  $|S| - n \in [2, 3]$ . It thus suffices to prove that

$$(26) \quad |\{0\} \cup \Sigma_{\leq |S| - n}(T)| \geq |S| - n + |\text{supp}(S)| - 1$$

in the two remaining cases. Let  $D = \{0\} \cup \text{supp}(T_1)$ . Since  $T_1$  is square-free and zero-sum free, we have

$$(27) \quad |D| = |T_1| + 1 \quad \text{and} \quad D \dot{+} D = \Sigma_{\leq 2}(T_1).$$

Since  $0 \notin \text{supp}(T)$  (per definition of  $T$ ) with  $T = T_0 T_1$ , we have  $0 \notin \text{supp}(T_0) \setminus \text{supp}(T_1)$ . Since  $T_1$  zero-sum free, we have  $0 \notin \Sigma_{\leq 2}(T_1)$ . Thus, in view of  $T = T_0 T_1$  and Claim 1, it follows that  $\text{supp}(T_0) \setminus \text{supp}(T_1)$  and  $\Sigma_{\leq 2}(T_1)$  are

both disjoint subsets of  $\Sigma_{\leq 2}(T)$  that do not contain 0. Combining this with (25) and (27), we obtain

$$\begin{aligned} |\{0\} \cup \Sigma_{\leq |S|-n}(T)| &\geq |\{0\} \cup \Sigma_{\leq 2}(T)| \geq 1 + |\text{supp}(T_0) \setminus \text{supp}(T_1)| \\ (28) \qquad \qquad \qquad &= 1 + |\text{supp}(T_0) \setminus \text{supp}(T_1)| \end{aligned}$$

It remains to estimate  $|D \dot{+} D|$  using Lemmas 2.8 and 2.9.

Suppose  $|S| - n = 2$ . Then, in view of (27) and (14), we have  $|D| = |T_1| + 1 \geq |S| - n + 2 = 4$ . If  $\text{supp}(T_1) \cup \{0\} = D = \langle D \rangle$  is an elementary 2 group, then  $0 \in \Sigma_3(T_1)$ , contradicting that  $T_1$  is zero-sum free. Therefore we may assume otherwise, in which case Lemma 2.8 and (27) together imply  $|D \dot{+} D| \geq |D| = |T_1| + 1 \geq |\text{supp}(T_1)| + 1$ . Applying this estimate in (28), and recalling that  $T = T_0 T_1$  with  $|\text{supp}(T)| \geq |\text{supp}(S)| - 1$ , we obtain

$$\begin{aligned} |\{0\} \cup \Sigma_{\leq |S|-n}(T)| &\geq 1 + |\text{supp}(T_0) \setminus \text{supp}(T_1)| + |\text{supp}(T_1)| \\ &= 2 + |\text{supp}(T)| \geq 1 + |\text{supp}(S)| = |\text{supp}(S)| \end{aligned}$$

Thus (26) is established in this case, as desired.

It remains to consider the case when  $|S| - n = 3$ . Then, in view of (27) and (14), we have  $|D| = |T_1| + 1 \geq |S| - n + 2 = 5$ . Let  $H = \langle D \rangle$ . If  $H$  is an elementary 2-group, then  $|D| \geq 5$  ensures that it must have size  $|H| \geq 8$ . Consequently, if  $|D| = |\text{supp}(T_1) \cup \{0\}| \geq |H| - 1$ , then it is easily seen that  $T_1$  will contain a 3-term zero-sum subsequence, contradicting that  $T_1$  is zero-sum free. On the other hand, if  $H$  is not an elementary 2-group and  $D = H$ , then there will be some  $a \in D \setminus \{0\} = \text{supp}(T_1)$  with  $\text{ord}(a) \geq 3$ . Since  $\{0\} \cup \text{supp}(T_1) = D = H$  ensures that we also have  $-a \in \text{supp}(T_1)$ , and since  $a \neq -a$  in view of  $\text{ord}(a) \geq 3$ , it follows that  $T_1$  contains a 2-term zero-sum, again contradicting that  $T_1$  is zero-sum free. Finally, since  $|D| \geq 5$ , we cannot have  $D = L \cup (a + L)$  with  $L \leq G$  an order 2 subgroup. As a result, Lemma 2.9 and (27) together imply  $|D \dot{+} D| \geq |D| + 1 = |T_1| + 2 \geq |\text{supp}(T_1)| + 2$ . Applying this estimate in (28), and recalling that  $T = T_0 T_1$  with

$|\text{supp}(T)| \geq |\text{supp}(S)| - 1$ , we obtain

$$\begin{aligned} |\{0\} \cup \Sigma_{\leq |S|-n}(T)| &\geq 1 + |\text{supp}(T_0) \setminus \text{supp}(T_1)| + |\text{supp}(T_1)| \\ &= 3 + |\text{supp}(T)| \geq 2 + |\text{supp}(S)| = |\text{supp}(T)| \end{aligned}$$

Thus (26) is established in the final case, completing the proof.  $\square$

#### 4. Concluding Remarks

Let  $G$  be a finite abelian group with exponent  $\exp(G)$ . Let  $S$  be a sequence over  $G$  with  $|S| \geq |G| + 1$  and  $0 \notin \Sigma_{|G|}(S)$ . When  $G$  is non-cyclic,  $|\text{supp}(S)| \leq |S| - |G| + 1$  and  $|S| \geq |G| + \exp(G) - 1$ , we can get better lower bounds for  $|\Sigma_{|G|}(S)|$  than those from Conjecture 1.2 (see Proposition 4.4). We need the following results.

**Proposition 4.1.** *(Gao and Leader, 2005) Let  $G$  be a finite abelian group and let  $S$  be a sequence over  $G$  with  $|S| \geq |G| + 1$  and  $0 \notin \Sigma_{|G|}(S)$ . Then there is a zero-sum free sequence  $T$  over  $G$  such that  $|T| = |S| - |G| + 1$  and  $|\Sigma_{|G|}(S)| \geq |\Sigma(T)|$ .*

For every integer  $k \in [1, D(G) - 1]$ , let  $f_G(k) = \min\{|\Sigma(T)| : T \in \mathcal{F}(G), |T| = k \text{ and } 0 \notin \Sigma(T)\}$

**Proposition 4.2.** *Let  $G$  be a finite abelian group that is noncyclic with exponent  $\exp(G)$ .*

- (1) *If  $k \geq \exp(G)$ , then  $f_G(k) \geq 2k - 1$ . (Olson and White, 1975; Sun, 2007)*

(2) If  $k \geq \exp(G) + 1$ , then  $f_G(k) \geq 3k - 1$ . (Gao, Li, Peng and Sun, 2008)

**Proposition 4.3.** (Pixton, 2009) *Let  $G$  be a finite abelian group and let  $T$  be a zero-sum free sequence over  $G$ .*

- (1) *If the rank of  $\langle \text{supp}(T) \rangle$  is at least 3, then  $|\Sigma(T)| \geq 4|T| - 5$ .*
- (2) *If the rank of  $\langle \text{supp}(T) \rangle$  is at least  $r$ , then  $|\Sigma(T)| \geq 2^r|T| - (r - 1)2^r - 1$ .*

Let  $G$  be a finite abelian group of rank  $r = r(G)$ . For every  $t \in [1, r]$ , define

$$d_t(G) = \max\{D(H) : H \leq G, r(H) = t\},$$

where the maximum is taken as  $H$  runs over all subgroups of  $G$  of rank  $t$ .

**Proposition 4.4.** *Let  $G$  be a finite abelian group that is noncyclic, let  $r = r(G)$  be the rank of  $G$ , and let  $S$  be sequence over  $G$  with  $|S| \geq |G| + 1$  and  $0 \notin \Sigma_{|G|}(S)$ .*

- (1) If  $|S| \geq |G| + \exp(G) - 1$ , then  $|\Sigma_{|G|}(S)| \geq 2|S| - 2|G| + 1$ .
- (2) If  $|S| \geq |G| + \exp(G)$ , then  $|\Sigma_{|G|}(S)| \geq 3|S| - 3|G| + 2$ .
- (3) If  $|S| \geq |G| + d_{t-1}(G) - 1$  with  $t \in [2, r]$ , then  $|\Sigma_{|G|}(S)| \geq 2^t|S| - 2^t|G| + (t-2)2^t - 1$ .
- (4) If  $|S| \geq |G| + d_2(G) - 1$ , then  $|\Sigma_{|G|}(S)| \geq 4|S| - 4|G| - 1$ .

*Proof.* We only prove Conclusion 3 here. The other three conclusions can be proved in a similar way. By Proposition 4.1, there is a zero-sum free sequence  $T$  over  $G$  with  $|T| = |S| - |G| + 1$  and  $|\Sigma_{|G|}(S)| \geq |\Sigma(T)|$ . Since  $|T| = |S| - |G| + 1 \geq d_{t-1}(G)$  and  $T$  is zero-sum free, the rank of  $\langle T \rangle$  is at least  $t$ . It follows from Proposition 4.3 that  $|\Sigma_{|G|}(S)| \geq |\Sigma(T)| \geq 2^t|T| - (t-1)2^t - 1 = 2^t(|S| - |G| + 1) - (t-1)2^t - 1 = 2^t|S| - 2^t|G| - (t-2)2^t - 1$ .  $\square$

Given a fixed (and arbitrary) finite abelian group  $G$ , it would be very difficult to give a sharp lower bound for  $|\Sigma_{|G|}(S)|$  involving  $|\text{supp}(S)|$  in general. Indeed, even finding sharp lower bounds when  $G$  is not fixed would be difficult, though it would be expected that the improvement be at least quadratic in  $|\text{supp}(S)|$ , rather than linear. We end this section with the following open problem.

**Conjecture 4.5.** *Let  $G$  be a finite abelian group and let  $S$  be a sequence over  $G$  with  $|S| \geq |G| + 1$  and  $0 \notin \Sigma_{|G|}(S)$ . Then there is a zero-sum free sequence  $T$  over  $G$  of length  $|T| = |S| - |G| + 1$  such that  $|\Sigma_{|G|}(S)| \geq |\Sigma(T)|$  and  $|\text{supp}(T)| \geq \min\{|S| - |G| + 1, |\text{supp}(S)| - 1\}$ .*

**Acknowledgements.** This work was supported by the National Key Basic Research Program of China (Grant No. 2013CB834204), the PC-SIRT Project of the Ministry of Science and Technology, the National Science Foundation of China, and the Education Department of Henan Province (Grant No. 2009A110012)

## References

- [1] B. Bollobás and I. Leader, *The number of  $k$ -sums modulo  $k$* , J. Number Theory, 78(1999) 27-35.
- [2] A. Bialostocki, D. Gryniewicz and M. Lotspeich *On some developments of the Erdős-Ginzburg-Ziv theorem. II*, Acta Arith., 110(2003) 173-184.
- [3] A. Bialostocki and M. Lotspeich, *Some developments of the Erdős-Ginzburg-Ziv theorem*, in "Sets, Graphs and Numbers", Coll. Math. Soc. J.Bolyai, 60(1992) 97-117.
- [4] J.D. Bovey, P. Erdős, I. Niven, *Conditions for zero sum modulo  $n$* , Canad. Math. Bull., 18(1975) 27-29.
- [5] Y. Caro, *Zero-sum problems- a survey*, Discrete Math., 152(1996) 93-113.
- [6] M. DeVos, L. Goddyn and B. Mohar, *A generalization of Kneser's Addition Theorem*, Adv. Math., 220(2009) 1531-1548.
- [7] R.B. Eggleton and P. Erdős, *two combinatorial problems in group theory*, Acta Arith., 21(1972) 111-116.
- [8] L. Gallardo, G. Grekos and J. Pinko, *On a variant of the Erdős-Ginzburg-Ziv problem*, Acta Arith., 89 (1999) 331-336.
- [9] W.D. Gao, *Addition theorems for finite abelian groups*, J. Number Theory, 53(1995) 241-246.
- [10] W.D. Gao, *A combinatorial problem on finite abelian groups*, J. Number Theory, 58(1996) 100-103.
- [11] W.D. Gao, *On the number of subsequences with given sum*, Discrete Mathematics, 195(1999) 127-138.
- [12] W.D. Gao and I. Leader, *Sums and  $k$ -sums in abelian groups of order  $k$* , J. Number Theory, 120(2006) 26-32.
- [13] W.D. Gao, Y.L. Li, J.T. Peng and F. Sun, *On subsequence sums of a zero-sum free sequence II*, Electron. J. Combin., 15 (2008), Research Paper 117, 21 pp.
- [14] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebra, Combinatorial and Analytic Theory, Pure and Applied Mathematics*, vol.278, Chapman & Hall/CRC, 2006.
- [15] D. Gryniewicz, *On a conjecture of Hamidoune for subsequence sums*, Integers, 5(2005) No.2, A7.
- [16] D. Gryniewicz, *Structural Additive Theory*, Developments in Mathematics, Springer, 2013.
- [17] D. J. Gryniewicz, E. Marchan, O. Ordaz, *Representation of finite abelian group elements by subsequence sums*, J. Théor. Nombres Bordeaux, 21 (2009), no. 3, 559–587.
- [18] Y. O. Hamidoune, *Subsequence sums*, Combinatorics, Probability and Computing, 12(2003) 413-425
- [19] H.B. Mann, *Two addition theorems*, J. Combinatorial Theory, 3(1967) 233-235.
- [20] M. B. Nathanson, *Additive Number Theory, Inverse Problems and the Geometry of Sumsets*, GMT 165, Springer-Verlag (New York, 1996).
- [21] J.E. Olson, *An addition theorem for finite abelian groups*, J. Number Theory 9(1977) 63-70.
- [22] J.E. Olson and E.T. White, *Sums from a sequence of group elements*, in: Number Theory and Algebra, Academic Press, New York, 1977, pp. 215-222.
- [23] A. Pixton, *Sequences with small subsum sets*, J. Number Theory, 129(2009) 806-817.
- [24] S. Savchev and F. Chen, *Long  $n$ -zero-free sequences in finite cyclic groups*, Discrete Mathematics, 308(2008) 1-8.
- [25] P. Scherk, *Distinct elements in a set of sums*, Amer. Math. Monthly, 62(1955) 46-47.
- [26] F. Sun, *On subsequence sums of a zero-sum free sequence*. Electron. J. Combin., 14 (2007), Research Paper 52, 9 pp.

# Thank you for your attention!

*E-mail address:* [wdgao@nankai.edu.cn](mailto:wdgao@nankai.edu.cn)